

Acknowledgments.—We thank Bradford Hollingsworth for providing the photo voucher number.

LITERATURE CITED

QUINTERO-DÍAZ, G. E., J. VÁZQUEZ-DÍAZ, AND J. J. SIGALA-RODRÍGUEZ. 2008. Reptiles. Pp. 141–145 *In* La Biodiversidad en Aguascalientes: Estudio de Estado. Comisión Nacional para el Conocimiento y Uso de la Biodiversidad (CONABIO), Instituto del Medio Ambiente del Estado de Aguascalientes (IMAE), Universidad Autónoma de Aguascalientes (UAA), Mexico.



Fig. 1. A *Hemidactylus frenatus* (SDSNH_HerpPC_05302) observed in Municipio de Valaparaíso, Zacatecas, Mexico.

📷 © Jorge A. Bañuelos Alamillo

JORGE A. BAÑUELOS-ALAMILLO¹, RUBÉN ALONSO CARBAJAL-MÁRQUEZ^{2,3}, AND ILSE YASARETH TRUJILLO DE LA TORRE⁴

¹Unidad Académica de Ciencias Biológicas, Universidad Autónoma de Zacatecas, Edificio de Biología Campus II Ave. Preparatoria S/N, Col. Agronómica, C. P. 98066, Zacatecas, Zacatecas, Mexico. E-mail: j.alberto.ba@gmail.com

²El Colegio de la Frontera Sur. Departamento de Conservación de la Biodiversidad. Unidad Chetumal, Av. Centenario Km 5.5, C. P. 77014, Chetumal, Quintana Roo, Mexico. E-mail: redman031@hotmail.com (Corresponding author)

³Conservación de la Biodiversidad del Centro de México, A. C. Andador Torre de Marfil No. 100, C. P. 20229, Aguascalientes, Aguascalientes, Mexico.

⁴Universidad Autónoma de Fresnillo, Carretera Federal Panamericana Km. 725.3 s/n Col. Lindavista, C. P. 99039, Fresnillo, Zacatecas, Mexico.

Family Gekkonidae

***Lepidodactylus lugubris* (Duméril & Bibron, 1836).** PANAMA: PROVINCIA DE BOCAS DEL TORO: Parque Nacional Marino Isla Bastimentos (9.315°N, 82.156°W; WGS 84); elev. 9 m; 12 November 2004; David Hegner. A photo voucher of this individual is deposited at the University of Texas at Arlington Digital Collection (UTADC-8629; Fig. 1A). The single individual of this introduced species was found active at night (ca. 1900 h) on the wall of old wooden house, a few meters away from primary lowland wet forest. The vicinity of the finding was anthropogenically modified, and several other wooden houses were present close to the bay at Bahía Honda (Fig. 1B), near the mangrove forest. This locality represents the first confirmation of this species from Isla Bastimentos, as previous information was unverified (see Sosa-Bartuano and Ramos-Reyes, 2015).

Hoogmoed and Avila-Pires (2015) provided a review of the distribution of *Lepidodactylus lugubris* in the New World, and for Panama included records from the literature and six localities for specimens from the province of Bocas del Toro in museum collections in the United States. The closest locality in Bocas del Toro is ca. 7 km from our record (Lotzkat, 2010), and thus the presence of this species on Isla Bastimentos was not surprising. Jiménez and Abarca (2015) provided several records for the closest localities in Costa Rica, from the province of Limón.



Fig. 1. (A) An individual of *Lepidodactylus lugubris* (UTADC-8629) from Isla Bastimentos, Provincia de Bocas del Toro, Panama; and (B) the location where it was found. © David Hegner

Acknowledgments.—We thank Carl J. Franklin for providing the photo voucher number.

LITERATURE CITED

- HOOGMOED, M. S., AND T. C. S. AVILA-PIRES. 2015. *Lepidodactylus lugubris* (Duméril & Bibron 1836) (Reptilia: Gekkonidae), an introduced lizard new for Brazil, with remarks on and correction of its distribution in the New World. *Zootaxa* 4,000: 90–110.
- JIMÉNEZ, R. R., AND J. G. ABARCA. 2015. Records for the Mourning Gecko (*Lepidodactylus lugubris*) and its expansion in Costa Rica. *Mesoamerican Herpetology* 2: 214–217.
- LOTZKAT, S. 2010. Reptiles de San San Pond Sak y áreas adyacentes en Bocas del Toro, Panama. Visual identification guides. 4 pp. (www.senckenberg.de/root/index.php?page_id=8958; accessed 7 January 2016).
- SOSA-BARTUANO, A., AND K. RAMOS-REYES. 2015. Distribution notes and comments on the predation of the Mourning Gecko, *Lepidodactylus lugubris* (Duméril & Bibron, 1836), in Panama. *Mesoamerican Herpetology* 2: 375–376.

DANIEL JABLONSKI¹, DAVID HEGNER², ÁNGEL SOSA-BARTUANO^{3,4}

¹Department of Zoology, Comenius University in Bratislava, Mlynská dolina B-1, 842 15, Bratislava, Slovakia.
E-mail: daniel.jablonski@balcanica.cz (Corresponding author)

²Mšenská 3938/26, 466 04 Jablonec nad Nisou, Czech Republic.

³Colaborador, Museo de Vertebrados de la Universidad de Panamá, Ciudad de Panamá, Panama.

⁴Red Mesoamericana y del Caribe para la Conservación de los Anfibios y Reptiles (Red MesoHerp).

Family: Iguanidae

***Iguana iguana* (Linnaeus, 1758).** Lee (1996: 210) noted the distribution of the wide-ranging Green Iguana (*Iguana iguana*) in the Yucatan Peninsula as “from southwestern Campeche, eastern Tabasco, and El Petén to Belize,” and documented specimens from Isla Cozumel, Quintana Roo, Mexico. Although Lee (1996; 2000) indicated the range of *I. iguana* in extreme southern Quintana Roo, near the Belize border, no specimens or specific localities were specified. Charruau et al. (2015) reported the presence of *I. iguana* on the Banco Chinchorro atoll, off the southeastern coast of Quintana Roo, near Belize. In this note I report three specimens of *I. iguana* from Quintana Roo that